

Phenomenological models of baryon

Do Quoc Tuan

Dept. of Theoretical Physics, College of Science, VNU, Hanoi, Vietnam

Abstract

In this paper, I present almost my works performed during my time at VNU. I was interested in the composite Skyrme model proposed by H. Y. Cheung and F. Gursey. The expanding for this model based on results obtained from the original Skyrme model given by T. H. R. Skyrme in 1961 was general and interesting.

Acknowledgments

I want to show deep gratitude to my supervisor Dr. Pham Thuc Tuyen (Department of Theoretical Physics, College of Science (another name is Hanoi University of Science), Vietnam National University) because of everything that he has been teaching and supporting me during a student's time!

I also thank Department of Theoretical Physics and Faculty of Physics for giving me the best conditions to completing the thesis!

Finally, and most of all, I want to give my thesis to my parent, my beloved as a special gift for the supports over a long period!

Ha noi, 6/2007
Do Quoc Tuan¹

¹email: do.tocxoan@gmail.com

Contents

1	Why is the Skyrme model	4
1.1	The weak point of QCD	4
1.2	The idea of Skyrme	5
2	The massless composite Skyrme model	9
2.1	The formalism	9
2.2	Calculations for static quantities of baryon	13
3	The composite Skyrme model with pion's mass	19
3.1	The term of pion'mass	19
3.2	The field equation and numerical results	20
4	The supersymmetric composite Skyrme model	23
5	Appendix	30
5.1	Some notations for Chapter 2 and 3	30
5.2	Some notations for Chapter 4	32

Chapter 1

Why is the Skyrme model

1.1 The weak point of QCD

In physically, we can divide the nature in two types of particles: matter particles and field particles. Matter particles interact together by fields concluding transfer interaction particles. There are four fields corresponding four interactions (four forces) and their particles. The graviton is a transfer interaction particle of gravitational force, the photon is a transfer interaction particle of electromagnetic force, $W^\pm$ and Z^0 are transfer interaction particle of weak force, the gluon is transfer interaction particle of strong force. It is known well, quarks are interact together by the strong force. In other word, the interaction of quarks is the exchange of gluons. And the theory that describes quarks and their interaction is *QCD* (**Q**uantum **C**hromo**D**ynamics). In mathematically, Gell-Mann and Neuman proposed that quarks make up a multiple which represents the gauge $SU(3)$ being the group of 3×3 unitary matrices with determinant **1**. Gell-Mann firstly gives generators of $SU(3)$, therefore, they are called Gell-Mann matrices λ . They satisfy the Lie algebra

$$[\lambda_i, \lambda_j] = 2if_{ijk}\lambda_k, \quad (1.1)$$

where f_{ijk} is the structure constant. It is totally antisymmetric

$$f_{ijk} = -f_{jik}. \quad (1.2)$$

To describe the interaction between quarks, we need using the gluon field A_μ^α . And now, the Lagrangian density is the Yang-Mills Lagrangian density

$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}^\alpha F^{\mu\nu\alpha} + i\bar{\psi}\gamma^\mu(\partial_\mu + igA_\mu)\psi + M\bar{\psi}\psi, \quad (1.3)$$

$$A_\mu = A_\mu^\alpha \lambda^\alpha, \quad (1.4)$$

$$F_{\mu\nu}^\alpha = \partial_\mu A_\nu^\alpha - \partial_\nu A_\mu^\alpha - gf^{\alpha\beta\gamma}A_\mu^\beta A_\nu^\gamma, \quad (1.5)$$

where $\mathbf{g}$ is a constant of interaction. At the high energy scale ($> TeV$), QCD is very good to use. But, when the energy lower (in GeV), it meets with some problems in which describe quantities of quarks and gluons. The main cause is the high value of $\mathbf{g}$, it obstructs an expansion of interaction terms of hadron. To repair this problem, t'Hooft proposed that $1/N_c$ plays as $\mathbf{g}$ (N_c is color number in a gauge theory). And, he gave that in large- N_c limit, QCD becomes equivalent to an effective field theory of meson.

1.2 The idea of Skyrme

In 1961, Skyrme gave the new model in which baryons may be obtained from the theory of meson [1]. He began with the Lagrangian density of pion

$$\mathcal{L} = \frac{1}{2} \partial_\mu \vec{\pi} \partial^\mu \vec{\pi} - \frac{m^2}{2} \vec{\pi} \vec{\pi}, \quad (1.6)$$

in which π 's make up a three dimensional isotopic space. After, he expanded the isotopic $SU(2)$ becomes a new group with the algebra

$$[V_i, V_j] = 2i\varepsilon_{ijk} V_k, \quad (1.7)$$

$$[A_i, A_j] = 2i\varepsilon_{ijk} V_k, \quad (1.8)$$

$$[V_i, A_j] = 2i\varepsilon_{ijk} A_k, \quad (1.9)$$

where generators V_i are vector charges, generators A_i are axial charges. And now, an isotopic space have four components $\{\vec{\pi}\} \rightarrow \{\vec{\varphi}, \varphi_4\}$ transforming in a rule

$$[V_i, \varphi_j] = 2i\varepsilon_{ijk} \varphi_k, \quad (1.10)$$

$$[V_i, \varphi_4] = 0, \quad (1.11)$$

$$[A_i, \varphi_j] = -2i\varepsilon_{ijk} \varphi_k, \quad (1.12)$$

$$[A_i, \varphi_4] = 2i\varphi_i. \quad (1.13)$$

Particles described by φ_i ($i = 1, 2, 3$) are pseudo- scalars and making isotopic vectors. A particle described by φ_4 has a positive parity, a spin and an isotopic spin equals zero, and it is a vacuum.

If replace V_i and A_i by L_i and R_i defined

$$L_i = \frac{1}{2} (V_i - A_i), \quad (1.14)$$

$$R_i = \frac{1}{2} (V_i + A_i), \quad (1.15)$$

then obtaining a new algebra

$$\begin{aligned}
[L_i, L_j] &= \left[\frac{1}{2} (V_i - A_i), \frac{1}{2} (V_j - A_j) \right], \\
&= \frac{1}{4} [V_i, V_j] - \frac{1}{4} [V_i, A_j] - \frac{1}{4} [A_i, V_j] + \frac{1}{4} [A_i, A_j], \\
&= \frac{1}{2} i \varepsilon_{ijk} V_k - \frac{1}{2} i \varepsilon_{ijk} A_k + \frac{1}{2} i \varepsilon_{jik} A_k + \frac{1}{2} i \varepsilon_{ijk} V_k, \\
&= i \varepsilon_{ijk} V_k - i \varepsilon_{jik} A_k, \\
&\Rightarrow [\mathbf{L}_i, \mathbf{L}_j] = 2\mathbf{i} \varepsilon_{ijk} \mathbf{L}_k
\end{aligned} \tag{1.16}$$

and

$$\begin{aligned}
[R_i, R_j] &= \left[\frac{1}{2} (V_i + A_i), \frac{1}{2} (V_j + A_j) \right], \\
&= \frac{1}{4} [V_i, V_j] + \frac{1}{4} [V_i, A_j] + \frac{1}{4} [A_i, V_j] + \frac{1}{4} [A_i, A_j], \\
&= \frac{1}{2} i \varepsilon_{ijk} V_k + \frac{1}{2} i \varepsilon_{ijk} A_k - \frac{1}{2} i \varepsilon_{jik} A_k + \frac{1}{2} i \varepsilon_{ijk} V_k, \\
&= i \varepsilon_{ijk} V_k + i \varepsilon_{jik} A_k, \\
&\Rightarrow [\mathbf{R}_i, \mathbf{R}_j] = 2\mathbf{i} \varepsilon_{ijk} \mathbf{R}_k
\end{aligned} \tag{1.17}$$

and

$$\begin{aligned}
[L_i, R_j] &= \left[\frac{1}{2} (V_i - A_i), \frac{1}{2} (V_j + A_j) \right], \\
&= \frac{1}{4} [V_i, V_j] + \frac{1}{4} [V_i, A_j] - \frac{1}{4} [A_i, V_j] - \frac{1}{4} [A_i, A_j], \\
&= \frac{1}{2} i \varepsilon_{ijk} V_k + \frac{1}{2} i \varepsilon_{ijk} A_k + \frac{1}{2} i \varepsilon_{jik} A_k - \frac{1}{2} i \varepsilon_{ijk} V_k, \\
&\Rightarrow [\mathbf{L}_i, \mathbf{R}_j] = \mathbf{0}.
\end{aligned} \tag{1.18}$$

This is an algebra of the chiral group $SU(2) \times SU(2)$. Thus, the expanding group is the chiral group $SU(2) \times SU(2)$. Continuously, he proposed that $\{\vec{\varphi}, \varphi_4\}$ form a three dimensional sphere

$$\vec{\varphi}^2 + \varphi_4^2 = 1. \tag{1.19}$$

Recently, $\{\vec{\varphi}, \varphi_4\}$ are associated with 2×2 unitary $U(x, t)$ matrix

$$\begin{aligned}
U(x, t) &= \exp i \vec{\tau} \vec{\pi}(x, t), \\
&= \exp i \vec{\tau} \frac{\vec{\pi}}{\pi} \pi,
\end{aligned}$$

$$= \cos \pi + i \vec{\tau} \frac{\vec{\pi}}{\pi} \sin \pi, \quad (1.20)$$

where $\pi = \sqrt{\vec{\pi}^2}$, τ 's are Pauli matrices. Pion fields now are associated with $\vec{\pi}$ fields

$$\varphi_4 \Leftrightarrow \cos \pi, \quad (1.21)$$

$$\vec{\varphi} \Leftrightarrow \frac{\vec{\pi}}{\pi} \sin \pi. \quad (1.22)$$

Really

$$\vec{\varphi}^2 + \varphi_4^2 = \frac{\vec{\pi}^2}{\pi^2} \sin^2 \pi + \cos^2 \pi = 1. \quad (1.23)$$

The Lagrangian density proposed by Skyrme [1] is

$$\mathcal{L} = \frac{F_\pi^2}{16} \text{Tr} (\partial_\mu U \partial_\mu U^\dagger) + \frac{1}{32e^2} \text{Tr} [(\partial_\mu U) U^\dagger, (\partial_\nu U) U^\dagger]^2, \quad (1.24)$$

here U is an $SU(2)$ matrix transforming as $U \rightarrow AUB^{-1}$ under an chiral $SU(2) \times SU(2)$. He proved that the model has a soliton solution (called hedgehog¹ solution)

$$U_o = \exp \left(i \vec{\tau} \frac{\vec{r}}{r} F(r) \right), \quad (1.25)$$

where $F(r)$ is the chiral angle, baryons are obtained from the quantization of soliton solution (see chapter 2).

It is well known that in the large- N limit, QCD may be regarded as the theory of effective meson fields. E. Witten argued that baryons may be regarded as solitons of this effective meson theory without an further reference to their quark content [2]. We know that solitons have not a spin. But, by quantizing around solitons, we obtain the special topological construct of these solitons which makes a spin. On the other hand, soliton solutions have a limited spatial distribution, therefore, they are correlative to particles that have a limited spatial dimension [2,15].

¹firstly called by Polyakov, see at **L. Ryder**, *Quantum Field Theory* (second edition), Cambridge University Press




Figure 1.1: The configuration of hedgehog, arrows on it are isotopic vectors of $\vec{c}$

Chapter 2

The massless composite Skyrme model

2.1 The formalism

In 1983, G. S. Adkins, C. R. Nappi and E. Witten used the Skyrme's idea to compute the static properties of baryons on the chiral group $SU(2) \times SU(2)$ and found that the model gives predictions that agree well with experiments [2]. However, this model has a problem in which the situation of particles at low energy are not interested but it regards that particles in free states, and this point can reduce the exactitude of theoretical datum. In 1989, this problem is treated by P. T. Tuyen and N. A. Viet [3]. They considered that at low energy particles are not free but they would shrink together into the cluster. Therefore, the group that describes them is $SU(2) \times SU(2) / SU(2)_{diag}$. This model gives few datum better than the sigma model. In 1990, H. Y. Cheung, F. Gursey [4] gave the general model in which the sigma model and the **V-T** model are two special cases corresponding with $n = 1$ and $n = 2$. They introduced a composite $SU(2)$ configuration $V_n = UPUP...PU$, where $U(x, t)$ is the $SU(2)$ matrix and $P = p_o + i\vec{\tau} \cdot \vec{p}$ is the constant $SU(2)$ matrix, and they proposed the composite Skyrme lagrangian density

$$\mathcal{L}_n = -\frac{F_\pi^2}{16n^2} Tr (\partial_\mu V^{-n} \partial^\mu V^n) + \frac{1}{32e^2 n^4} Tr \left([V^{-n} \partial_\mu V^n, V^{-n} \partial_\nu V^n]^2 \right), \quad (2.1)$$

with F_π is the pion decay constant, e is a dimensionless parameter.

Under the chiral group $SU(2) \times SU(2)$, U transforms as $U \rightarrow AUB^\dagger$, P transforms as $P \rightarrow BPA^\dagger$, and so V_n transforms as $V_n \rightarrow AV_n B^\dagger$. In the limit $P \rightarrow I$, $V_n \rightarrow U_n$, the lagrangian (2.1) becomes approximated by the lagrangian density

$$\mathcal{L}_n = -\frac{F_\pi^2}{16n^2} Tr (\partial_\mu U^{-n} \partial^\mu U^n) + \frac{1}{32e^2 n^4} Tr \left([U^{-n} \partial_\mu U^n, U^{-n} \partial_\nu U^n]^2 \right). \quad (2.2)$$

Now, using the hedgehog soliton solution gave by Skyrme $U_0(r) = \exp[i\tau \cdot \hat{r} F(r)]$, where τ 's are Pauli's matrices and $\hat{r} = \vec{r}/r$, the lagrangian density (2.2) is

$$\mathcal{L}_n = \frac{e^2 F_\pi^4}{8} \left[\left(\frac{dF}{d\tilde{r}} \right)^2 + 2 \frac{\sin^2 nF}{n^2 \tilde{r}^2} \right] + \frac{e^2 F_\pi^4 \sin^2 nF}{2n^2 \tilde{r}^2} \left[2 \left(\frac{dF}{d\tilde{r}} \right)^2 + \frac{\sin^2 nF}{n^2 \tilde{r}^2} \right], \quad (2.3)$$

where $\tilde{r} = eF_\pi r$ is the dimensionless variable.

And the static energy of hedgehog is

$$\mathcal{E}_n = \frac{F_\pi}{e} A_n = M_n, \quad (2.4)$$

$$A_n = \int_0^\infty 4\pi \tilde{r}^2 \left\{ \frac{1}{8} \left[\left(\frac{dF}{d\tilde{r}} \right)^2 + 2 \frac{\sin^2 nF}{n^2 \tilde{r}^2} \right] + \frac{e^2 F_\pi^4 \sin^2 nF}{2n^2 \tilde{r}^2} \left[2 \left(\frac{dF}{d\tilde{r}} \right)^2 + \frac{\sin^2 nF}{n^2 \tilde{r}^2} \right] \right\} d\tilde{r}. \quad (2.5)$$

We obtain the non-linear differential equation of $F(r)$ from minimum condition of the hedgehog's energy

$$\delta_F \mathcal{E}_n = 0, \quad (2.6)$$

or

$$\frac{d}{dr} \frac{\delta \mathcal{E}_n}{\delta F'} - \frac{\delta \mathcal{E}_n}{\delta F} = 0, \quad (Euler - Lagrange - equation) \quad (2.7)$$

$$\left(\frac{\tilde{r}^2}{4} + \frac{2}{n^2} \sin^2 nF \right) F'' + \frac{\tilde{r}}{2} F' + \frac{1}{n} \sin 2nF \left(F'^2 - \frac{\sin^2 nF}{n^2 \tilde{r}^2} - \frac{1}{4} \right) = 0. \quad (2.8)$$

To solve this equation we need boundary conditions for the $F(r)$. The first condition of $F(r)$ that relates to the limit of hedgehog's energy when $r \rightarrow \infty$ is $F(\infty) = 0$. The second condition relating to the exist of baryonic number B is $F(0) = \pi$. This equation can not give us the analytic solution. So we will solve it by the numerical method.

Now we will quantize the soliton by collective coordinates $A(t) = a_o(t) + i\vec{a}(t) \cdot \vec{\tau}$ with $a_\mu(t)$ ($\mu = 0, 1, 2, 3$) are collective coordinates, $A(t)$ is $SU(2)$ matrix. From properties of $SU(2)$ matrix, we have

$$A(t) A^\dagger(t) = 1,$$

$$\{a_o(t) + i\vec{a}(t) \cdot \vec{\tau}\} \{a_o(t) - i\vec{a}(t) \cdot \vec{\tau}\} = 1,$$

$$\mathbf{a}_\mu \mathbf{a}_\mu = \mathbf{1}. \quad (2.9)$$

By time derivative, resulting

$$\mathbf{a}_\mu \dot{\mathbf{a}}_\mu = \mathbf{0} \quad (2.10)$$




Figure 2.1: The numerical solution of (2.8) for cases of $n=2$ and $n=4$ [22, 25].

By this way, solitons to be particles. Different quantization take different particles. Thus, it takes us to the new idea about the origin of particles in nature in which they are made by a basic particle (called soliton). With the $A(t)$ matrix, the $U^n(x, t)$ will transform as

$$U^n(x, t) = A(t) U_0^n(x) A^{-1}(t)$$

, and we put it into (2.2), the lagrangian now is

$$\begin{aligned} L_n &= -\mathcal{E}_n + \Gamma_n \text{Tr} [\partial_0 A(t) \partial_0 A^{-1}(t)] , \\ &= -\mathcal{E}_n + 2\Gamma_n \sum_{\mu=0}^3 \dot{a}_\mu^2, \end{aligned} \quad (2.11)$$

with

$$\Gamma_n = \frac{1}{e^3 F_\pi} \Phi_n, \quad (2.12)$$

$$\Phi_n = \frac{2\pi}{3n^2} \int_0^\infty \tilde{r}^2 \sin^2 nF \left\{ 1 + 4 \left[\left(\frac{dF}{d\tilde{r}} \right)^2 + \frac{\sin^2 nF}{n^2 \tilde{r}^2} \right] \right\} d\tilde{r}. \quad (2.13)$$

The conjugate momenta are obtained from the derivative of lagrangian following $\dot{a}_\mu(t)$

$$\pi_\mu = \frac{\partial L_n}{\partial \dot{a}_\mu},$$

$$\begin{aligned}
&= \frac{\partial \left[-M_n + 2\Gamma_n \sum_{\mu=0}^3 \dot{a}_\mu^2 \right]}{\partial \dot{a}_\mu}, \\
&= 4\Gamma_n \dot{a}_\mu.
\end{aligned} \tag{2.14}$$

We go to the hamiltonian

$$\begin{aligned}
H_n &= \pi_\mu \dot{a}_\mu - L_n, \\
&= 4\Gamma_n \dot{a}_\mu \dot{a}_\mu - L_n, \\
&= M_n + 2\Gamma_n \dot{a}_\mu \dot{a}_\mu, \\
&= M_n + \frac{\left(\sum_{\mu=0}^3 \pi_\mu^2 \right)}{8\Gamma_n}.
\end{aligned} \tag{2.15}$$

Canonical quantizing momenta $\pi_\mu = -i \frac{\partial}{\partial a_\mu}$ let the hamiltonian as

$$H_n = M_n + \frac{1}{8\Gamma_n} \sum_{\mu=0}^3 \left(\frac{-\partial^2}{\partial a_\mu^2} \right). \tag{2.16}$$

Now, we will define the isotopic vector $\vec{c}$ following the way

$$\begin{aligned}
\dot{A}^\dagger A &= (\dot{a}_0 - i\dot{a}_i \tau_i) (a_0 + ia_i \tau_i), \\
&= \dot{a}_0 a_0 + \dot{\vec{a}} \vec{a} - i \left(a_0 \dot{\vec{a}} - \dot{a}_0 \vec{a} + \vec{a} \times \dot{\vec{a}} \right) \vec{\tau},
\end{aligned}$$

note (2.10), we have

$$\dot{A}^\dagger A = -i \vec{c} \vec{\tau}, \tag{2.17}$$

with

$$\vec{c} = \left(a_0 \dot{\vec{a}} - \dot{a}_0 \vec{a} + \vec{a} \times \dot{\vec{a}} \right). \tag{2.18}$$

The square isotopic vector $\vec{c}$ is

$$\begin{aligned}
\vec{c}^2 &= \dot{a}_0^2 \vec{a}^2 + \dot{\vec{a}}^2 a_0^2 + \vec{a}^2 \dot{\vec{a}}^2 - (\vec{a} \dot{\vec{a}})^2 - 2\dot{a}_0 a_0 \dot{\vec{a}} \vec{a}, \\
&= \dot{a}_0^2 \vec{a}^2 + \dot{\vec{a}}^2 a_0^2 + \vec{a}^2 \dot{\vec{a}}^2 - (\vec{a} \dot{\vec{a}})^2 + 2(\vec{a} \dot{\vec{a}})^2,
\end{aligned}$$

$$= \dot{a}_\mu \dot{a}_\mu. \quad (2.19)$$

In the configuration of hedgehog, since the isotopic vector and the radius vector have the same direction, rotation in the co-ordinate space can make an isospin of particle, and we assume that a spin equals to an isospin. The canonical momenta that correspond with collective coordinates are isospins. We can homogenize $\vec{c}^2$ with the square of a spin and an isospin

$$\vec{c}^2 = \frac{1}{16\Gamma_n^2} \vec{J}^2 = \frac{1}{16\Gamma_n^2} \vec{T}^2. \quad (2.20)$$

Thus, we have

$$H_n = M_n + \frac{1}{8\Gamma_n} \vec{J}^2 = M_n + \frac{1}{8\Gamma_n} \vec{T}^2, \quad (2.21)$$

with $\vec{J}$ is a spin and $\vec{T}$ is an isospin. Eigenvalues of hamiltonian (2.21) are

$$E_n = M_n + \frac{l(l+2)}{8\Gamma_n}, \quad (2.22)$$

with $l=2J$ and J is the quantum number of spin. So, the nucleon ($J = 1/2$) and the delta ($J = 3/2$) masses are given by

$$M_N = M_n + \frac{3}{8\Gamma_n}, \quad (2.23)$$

$$M_\Delta = M_n + \frac{15}{8\Gamma_n}. \quad (2.24)$$

Note (2.12), (2.13), (2.23), (2.24) we have

$$F_\pi = \sqrt[4]{\frac{(5M_N - M_\Delta)^3 (M_\Delta - M_N) \Phi_n}{384A_n^3}}, \quad (2.25)$$

$$e = \frac{4A_n F_\pi}{(5M_N - M_\Delta)}. \quad (2.26)$$

2.2 Calculations for static quantities of baryon

After constructing the formalism of massless composite model, we apply to the baryon (nucleon). Since this model is constructed depend on some ideas of theory of topology then we can start at the topological current to calculations for static quantities of baryon [2, 3, 4, 15].

The topological current (a baryonic current) is defined

$$B^\mu = \frac{\varepsilon^{\mu\nu\alpha\beta}}{24\pi^2 n} \text{Tr} \left[(U^{\dagger n} \partial_\nu U^n) (U^{\dagger n} \partial_\alpha U^n) (U^{\dagger n} \partial_\beta U^n) \right], \quad (2.27)$$

with $\varepsilon_{0123} = -\varepsilon^{0123} = 1$ (totally antisymmetry).

If we put $U^n(x, t) = A(t) U_0^n(x) A^{-1}(t)$ into (2.27), we will have

$$B^0 = -\frac{1}{2\pi^2} \frac{\sin^2 nF}{r^2} F', \quad (2.28)$$

$$B^i = \frac{i\varepsilon^{ijk} \sin^2 nF}{2\pi^2 r^2} F' x_k \text{Tr} \left[\dot{A}^\dagger A \tau_j \right]. \quad (2.29)$$

Note (2.17) and (2.20), then (2.29) is now

$$B^i = \frac{\varepsilon^{ijk} \sin^2 nF}{2\pi^2 r^2} F' x_k \frac{J_j}{2\Gamma_n}. \quad (2.30)$$

The baryonic number is

$$B = \int_0^\infty 4\pi r^2 B^0(r) dr = 1. \quad (2.31)$$

The iso-scalar electric square radius is

$$\langle r^2 \rangle_{E, I=0} = \int_0^\infty 4\pi r^2 B^0(r) r^2 dr. \quad (2.32)$$

The iso-scalar magnetic square radius is

$$\langle r^2 \rangle_{M, I=0} = \frac{\int_0^\infty 4\pi r^2 B^0(r) r^4}{5 \langle r^2 \rangle_{E, I=0}}. \quad (2.33)$$

We have the time's component of vector current $\vec{V}$

$$V_i^0 = \frac{i \sin^2 nF}{3 n^2 r^2} \left(F_\pi^2 + \frac{4}{e^2} \left[\left(\frac{dF}{dr} \right)^2 + \frac{\sin^2 nF}{n^2 r^2} \right] \right) \text{Tr} \left(\dot{A} A^\dagger \tau_i \right), \quad (2.34)$$

note (2.17)

$$V_i^0 = \frac{\sin^2 nF}{3n^2 r^2} \left(F_\pi^2 + \frac{4}{e^2} \left[\left(\frac{dF}{dr} \right)^2 + \frac{\sin^2 nF}{n^2 r^2} \right] \right) \frac{T_i}{2\Gamma_n}. \quad (2.35)$$

The density of charge of nucleon is defined by Gell-Mann - Nishijima

$$Q = \int_0^\infty \rho_{nucleon}(r) dr = I_3 + \frac{1}{2}B, \quad (2.36)$$

and

$$I_3 = \int_0^\infty r^2 V_3^0 d\Omega dr = \int_0^\infty \frac{4\pi \sin^2 nF}{3n^2} \left(F_\pi^2 + \frac{4}{e^2} \left[\left(\frac{dF}{dr} \right)^2 + \frac{\sin^2 nF}{n^2 r^2} \right] \right) \frac{T_3}{2\Gamma_n} dr. \quad (2.37)$$

So, we have

$$\rho_{nu}(r) = \frac{4\pi \sin^2 nF}{3n^2} \left(F_\pi^2 + \frac{4}{e^2} \left[\left(\frac{dF}{dr} \right)^2 + \frac{\sin^2 nF}{n^2 r^2} \right] \right) \frac{T_3}{2\Gamma_n} - \frac{\sin^2 nF}{\pi} F'. \quad (2.38)$$

With $\mathbf{T}_3 = \frac{1}{2}$ and $\mathbf{T}_3 = -\frac{1}{2}$, we have the charge's densities of proton and neutron.

After integrating, we obtain charges of a proton and a neutron

$$Q_{pro} = \int_0^\infty \rho_{pro}(r) dr = 1, \quad (2.39)$$

$$Q_{neu} = \int_0^\infty \rho_{neu}(r) dr = 0. \quad (2.40)$$

The iso-vector electric square radius and the iso-vector magnetic square radius are

$$\langle r^2 \rangle_{M, I=1} = \langle r^2 \rangle_{E, I=1} = \int_0^\infty r^2 V_3^0 dr = \infty. \quad (2.41)$$

However, in the experiment the iso-vector electric radius is **0.88**, the iso-vector magnetic radius is **0.8**. This is the weak point of the massless composite Skyrme model!

The iso-scalar magnetic moment is defined

$$\vec{\mu}_{I=0} = \frac{1}{2} \int (\vec{r} \times \vec{B}) d^3x, \quad (2.42)$$

and the result is

$$\vec{\mu}_{I=0} = \frac{1}{3\Gamma_n} \langle r^2 \rangle_{I=0} \vec{J}. \quad (2.43)$$

The connection between the spin and the magnetic moment is

$$\vec{\mu} = \frac{g}{2M_N} \vec{J}. \quad (2.44)$$

Note (2.23, 2.24), we have

$$g_{I=0} = g_{proton} + g_{neutron} = \frac{4}{9} M_N (M_\Delta - M_N) \langle r^2 \rangle_{I=0}. \quad (2.45)$$

The iso-vector magnetic moment is defined

$$\vec{\mu}_{I=1} = \frac{1}{2} \int (\vec{r} \times \vec{V}_3) d^3x, \quad (2.46)$$

where $\vec{V}_3$ is the vector current. After few calculations, the result is

$$\vec{\mu}_{I=1} = \frac{1}{3} \Gamma_n \vec{J}. \quad (2.47)$$

And we also have

$$g_{I=1} = g_{proton} - g_{neutron} = \frac{2M_N}{M_\Delta - M_N}. \quad (2.48)$$

Using (2.44) and (2.47), we can find

$$\mu_{pro} = \frac{g_{pro}}{2} = \frac{1}{4} \left[\frac{4}{9} M_N (M_\Delta - M_N) \langle r^2 \rangle_{I=0} + \frac{2M_N}{M_\Delta - M_N} \right], \quad (2.49)$$

$$\mu_{neu} = \frac{g_{neu}}{2} = \frac{1}{4} \left[\frac{4}{9} M_N (M_\Delta - M_N) \langle r^2 \rangle_{I=0} - \frac{2M_N}{M_\Delta - M_N} \right]. \quad (2.50)$$

The axial coupling is defined

$$g_A = -\frac{\pi}{3e^2} D_n, \quad (2.51)$$

with

$$D_n = \int_0^\infty \tilde{r}^2 \left\{ \frac{\sin 2nF}{n\tilde{r}} \left[1 + 4 \left(F'^2 + \frac{\sin^2 nF}{n^2 \tilde{r}^2} \right) \right] + F' \left(1 + \frac{8 \sin^2 nF}{n^2 \tilde{r}^2} \right) \right\} d\tilde{r}. \quad (2.52)$$




Figure 2.2: Plots of density of neutron charge and proton charge for a case of $n = 2$ [25].




Figure 2.3: Plots of density of neutron charge and proton charge for a case of $n = 4$ [25].

NUMERICAL RESULTS TABLE FOR THE CASE OF N=4 [25]

Quantity	Prediction	Experiment
e	1.10	-
$F_\pi(\text{MeV})$	168	186
$\langle r^2 \rangle_{E,I=0}^{1/2}(\text{fm})$	1.10	0.72
$\langle r^2 \rangle_{M,I=0}^{1/2}(\text{fm})$	1.93	0.81
$\langle r^2 \rangle_{E,I=1}^{1/2}(\text{fm})$	∞	0.88
$\langle r^2 \rangle_{M,I=1}^{1/2}(\text{fm})$	∞	0.8
$\mu_{pro}(\text{mag})$	2.55	2.79
$\mu_{neu}(\text{mag})$	-0.66	-1.91
$\left \frac{\mu_{pro}}{\mu_{neu}} \right $	3.86	1.46
g_A	3.52	1.23

Chapter 3

The composite Skyrme model with pion's mass

3.1 The term of pion's mass

When Skyrme constructed a model from the theory of pi-meson, he did not let the term of mass of pion because of its spontaneous symmetry breaking. However, with $n = 1, 2, 3, 4$, datum of prediction are not really exact. One of causes that we do not let the term of pion's mass into the lagrangian [2, 3]. Therefore, we will propose the composite lagrangian density with the pion's mass [22, 24].

The lagrangian density now is

$$\mathcal{L}_n^{pion} = \mathcal{L}_n + \frac{1}{8n} m_\pi^2 F_\pi^2 [Tr(U^n) - 2]. \quad (3.1)$$

Now using the hedgehog form gave by Skyrme $U_o(r) = \exp[i\tau \cdot \hat{r} F(r)]$, where τ 's are Pauli's matrices and $\hat{r} = \frac{\vec{r}}{|\vec{r}|}$, the lagrangian density is

$$\mathcal{L}_n^{pion} = \mathcal{L}_n + \frac{1}{4n} m_\pi^2 F_\pi^2 (\cos nF - 1), \quad (3.2)$$

$$\mathcal{L}_n = \frac{F_\pi^2}{8} \left[\left(\frac{dF}{dr} \right)^2 + 2 \frac{\sin^2 nF}{n^2 r^2} \right] + \frac{\sin^2 nF}{2e^2 n^2 r^2} \left[2 \left(\frac{dF}{dr} \right)^2 + \frac{\sin^2 nF}{n^2 r^2} \right]. \quad (3.3)$$

The energy of hedgehog is

$$\mathcal{E}_n^{pion} = \frac{F_\pi}{e} A_n + \frac{1}{e^3 F_\pi} \Theta_n, \quad (3.4)$$

with

$$A_n = \int_0^\infty 4\pi \tilde{r}^2 \left\{ \left[\frac{F'^2}{8} + \frac{\sin^2 nF}{4n^2 \tilde{r}^2} \right] + \frac{e^2 F_\pi^4 \sin^2 nF}{n^2 \tilde{r}^2} \left[F'^2 + \frac{\sin^2 nF}{2n^2 \tilde{r}^2} \right] \right\} d\tilde{r}, \quad (3.5)$$

and

$$\Theta_n = \int_0^\infty \frac{\pi \tilde{r}^2 m_\pi^2}{n} (\cos nF - 1) d\tilde{r}, \quad (3.6)$$

$\tilde{r} = eF_\pi r$ is a dimensionless variable.

After some steps of calculation as **chapter 2**, we go to the hamiltonian

$$H_n = \mathcal{E}_n^{pion} + \frac{1}{8\Gamma_n} \vec{J}^2 = \mathcal{E}_n^{pion} + \frac{1}{8\Gamma_n} \vec{T}^2, \quad (3.7)$$

with $\vec{J}$ is a spin and $\vec{T}$ is an isospin. Eigenvalues of hamiltonian (3.7) are

$$M_n = E_n = \frac{F_\pi}{e} A_n + \frac{1}{e^3 F_\pi} \Theta_n + \frac{l(l+2)}{8\Gamma_n}, \quad (3.8)$$

$l = 2J$ and J is the quantum number of spin.

So, the nucleon ($J = 1/2$) and delta ($J = 3/2$) masses are given by

$$M_N = \frac{F_\pi}{e} A_n + \frac{1}{e^3 F_\pi} \Theta_n + \frac{3e^3 F_\pi}{8\Phi_n}, \quad (3.9)$$

$$M_\Delta = \frac{F_\pi}{e} A_n + \frac{1}{e^3 F_\pi} \Theta_n + \frac{15e^3 F_\pi}{8\Phi_n}. \quad (3.10)$$

From (3.9) and (3.10), we obtain

$$F_\pi = \frac{2(M_\Delta - M_N) \Phi_n}{3e^3}, \quad (3.11)$$

$$e = \sqrt[4]{\frac{4(M_\Delta - M_N)^3 \Phi_n^2 A_n}{6M_N (M_\Delta - M_N)^2 \Phi_n - 9\Theta_n - 3(M_\Delta - M_N)^2 \Phi_n}}. \quad (3.12)$$

3.2 The field equation and numerical results

We obtain the nonlinear differential equation of $F(r)$ from minimum condition of the hedgehog's energy

$$\delta_F \mathcal{E}_n^{pion} = 0, \quad (3.13)$$

or

$$\frac{d}{dr} \frac{\delta \mathcal{E}_n^{pion}}{\delta F'} - \frac{\delta \mathcal{E}_n^{pion}}{\delta F} = 0 \text{ (Euler - Lagrange - equation)}, \quad (3.14)$$

$$\left(\frac{\tilde{r}^2}{4} + \frac{2}{n^2} \sin^2 nF \right) F'' + \frac{\tilde{r}}{2} F' + \frac{\sin 2nF}{n} \left(F'^2 - \frac{\sin^2 nF}{n^2 \tilde{r}^2} - \frac{1}{4} \right) - \beta \tilde{r}^2 \sin nF = 0, \quad (3.15)$$

where

$$\beta = \frac{m_\pi^2}{4e^2 F_\pi^2}. \quad (3.16)$$

We see that (3.16) contains the constant of e and F_π but they are not known because they are must found after solving (3.15). Thus, we will solve (3.15) by the way is taking the value of β in spite of e and F_π and raising it until it equals approximately the β that defined by (3.11, 3.12, 3.16). In this case $n = 4$, we need the value of β is 0.0025 [24].




Figure 3.1: Plots of numerical solution for a case of $n = 4$ [24].




Figure 3.2: Plots of density of neutron charge and proton charge for a case of $n = 4$ [24].

Following results that are obtained above by us (**chapter 2**), quantities of baryon (proton and neutron) are described below [24]

NUMERICAL RESULTS TABLE FOR THE CASE OF N=4

Quantity	Prediction	Experiment
e	2.87	-
$F_\pi(\text{MeV})$	446.56	186
$\langle r^2 \rangle_{E,I=0}^{1/2}(\text{fm})$	0.21	0.71
$\langle r^2 \rangle_{E,I=1}^{1/2}(\text{fm})$	2.26	0.88
$\langle r^2 \rangle_{M,I=0}^{1/2}(\text{fm})$	1.0	0.81
$\langle r^2 \rangle_{M,I=1}^{1/2}(\text{fm})$	2.26	0.8
$\mu_{pro}(\text{mag})$	1.64	2.79
$\mu_{neu}(\text{mag})$	-1.57	-1.91
$\left \frac{\mu_{pro}}{\mu_{neu}} \right $	1.04	1.46
g_A	1.22	1.23

Chapter 4

The supersymmetric composite Skyrme model

The fundamental theories of particles consider that elementary particles are points. They all have the spin half-one and combining them together will make particles which have the structural space. But, the model gave by Skyrme is quite otherwise. He consider that there are solitons. In these models, one consider that soliton solutions have the structural space and the spin zero. By quantizing these solitons, we obtain real particles that have the spin non-zero. Since working with solitons, these models are all nonlinear. There is an important point that these models work at the ideal case (a pion's mass is zero). That makes datum of prediction are not really exact. Thus, we need to let the term of pion's mass in these models. But, with only pion, the models would not be exact, because we also have other mesons. Must let all of mesons into the model. Thus, we extend the model onto the $SU(3)$ group. But then will be an another cause that makes the model is not really exact. In general we must consider the contributions of the particle partners of the mesons. On the other hand, each meson has a "superpartner" (make up from quark superpartners). When all of these contributions are counted in the model, it may be exact [23].

In 1984, E. A. Bergshoeff, R. I. Nepomachie, H. J. Schnitzer proposed Skyrmion of four-dimensional supersymmetric non-linear sigma model [5]. The composite Skyrme model proposed by H. Y. Cheung and F. Gursey in 1990. It's the general model in which the sigma model and $V - T$ model are two special cases corresponding $n = 1, 2$. Thus, we will calculate the composite model in the supersymmetry by the way proposed by E. A. Bergshoeff et al [26].

We recall the Lagrangian density of composite Skyrme model

$$\mathcal{L}_n = -\frac{f_\pi^2}{16n^2} \text{Tr} (\partial_\mu U^{-n} \partial^\mu U^n) + \frac{1}{32e^2 n^4} \text{Tr} ([U^{-n} \partial_\mu U^n, U^{-n} \partial_\nu U^n]^2), \quad (4.1)$$

with f_π is the pion decay constant, e is a dimensionless parameter.

Replacing ordinary derivatives in the Lagrangian density into covariant derivatives

$$D_\mu U^n = \partial_\mu U^n - iV_\mu U^n \tau_3, \quad (4.2)$$

the Lagrangian density (4.1) is now

$$\mathcal{L}_n = -\frac{f_\pi^2}{16n^2} \text{Tr} (D_\mu U^{-n} D^\mu U^n) + \frac{1}{32e^2 n^4} \text{Tr} \left([U^{-n} D_\mu U^n, U^{-n} D_\nu U^n]^2 \right). \quad (4.3)$$

and is invariant under local $U(1)_R$ and global $SU(2)_L$ transformations

$$U^n(r) \rightarrow AU^n(x) e^{i\lambda(r)\tau_3}, A \in SU(2)_L, \quad (4.4)$$

$$V_\mu(r) \rightarrow V_\mu(r) + \partial_\mu \lambda(r). \quad (4.5)$$

Really, we have

$$\begin{aligned} (D_\mu U^{-n})' &= [\partial_\mu + i(V_\mu + \partial_\mu \lambda) \tau_3] (U^{-n} e^{-i\lambda(x)\tau_3}) A^\dagger, \\ &= (\partial_\mu U^{-n}) e^{-i\lambda\tau_3} A^\dagger - i(\partial_\mu \lambda) \tau_3 U^{-n} e^{-i\lambda\tau_3} A^\dagger + \\ &\quad + iV_\mu \tau_3 U^{-n} e^{-i\lambda\tau_3} A^\dagger + i(\partial_\mu \lambda) \tau_3 U^{-n} e^{-i\lambda\tau_3} A^\dagger, \\ &= [(\partial_\mu + iV_\mu \tau_3) U^{-n}] e^{-\lambda(x)\tau_3} A^\dagger, \\ &= (\mathbf{D}_\mu \mathbf{U}^{-\mathbf{n}}) \mathbf{e}^{-i\lambda(\mathbf{x})\tau_3} \mathbf{A}^\dagger; \end{aligned} \quad (4.6)$$

$$\begin{aligned} (D^\mu U^n)' &= A [\partial^\mu - i(V^\mu + \partial^\mu \lambda) \tau_3] (U^n e^{i\lambda(x)\tau_3}), \\ &= A (\partial^\mu U^n) e^{i\lambda\tau_3} + iA (\partial^\mu \lambda) \tau_3 U^n e^{i\lambda\tau_3} - iAV^\mu \tau_3 U^n e^{i\lambda\tau_3} - iA (\partial^\mu \lambda) \tau_3 U^n e^{i\lambda\tau_3}, \\ &= A (\partial^\mu U^n) e^{i\lambda\tau_3} - iAV^\mu \tau_3 U^n e^{i\lambda\tau_3} = A [(\partial^\mu - iV^\mu \tau_3) U^n] e^{i\lambda\tau_3}, \\ &= \mathbf{A} (\mathbf{D}^\mu \mathbf{U}^{\mathbf{n}}) \mathbf{e}^{i\lambda\tau_3}; \end{aligned} \quad (4.7)$$

$$\Rightarrow (D_\mu U^{-n})' (D^\mu U^n)' = (D_\mu U^{-n}) e^{-i\lambda(x)\tau_3} A^\dagger A e^{i\lambda\tau_3} (D^\mu U^n),$$

$$= (D_\mu U^{-n}) (D^\mu U^n). \quad (4.8)$$

$$\Rightarrow \text{Tr} \left[(D_\mu U^{-n})' (D^\mu U^n)' \right] = \text{Tr} \left[(D_\mu U^{-n}) (D^\mu U^n) \right]. \quad (4.9)$$

The second term of Lagrangian is proved similarly.

The gauge field $V_\mu(r)$ is defined

$$V_\mu = -\frac{i}{2n} \text{Tr} (U^{-n} \partial_\mu U^n \tau_3). \quad (4.10)$$

We parameterize the $SU(2)$ matrix $U^n(r)$ in terms of the complex scalars A_i

$$U^n(r) = \begin{pmatrix} A_1 & -A_2^* \\ A_2 & A_1^* \end{pmatrix}, \quad (4.11)$$

with the unitary constraint

$$U^\dagger U = 1 \Rightarrow \bar{A}^i A_i = A_1^* A_1 + A_2^* A_2 = 1. \quad (4.12)$$

Now, (4.2) is

$$D_\mu \begin{pmatrix} A_1 & -A_2^* \\ A_2 & A_1^* \end{pmatrix} = \partial_\mu \begin{pmatrix} A_1 & -A_2^* \\ A_2 & A_1^* \end{pmatrix} - iV_\mu \begin{pmatrix} A_1 & A_2^* \\ A_2 & -A_1^* \end{pmatrix} \quad (4.13)$$

or

$$D_\mu A_i = (\partial_\mu - iV_\mu) A_i, \quad (4.14)$$

$$D_\mu \bar{A}_i = (\partial_\mu + iV_\mu) \bar{A}_i. \quad (4.15)$$

The form of gauge field is then

$$\begin{aligned} V_\mu &= -\frac{i}{2n} \text{Tr} (U^{-n} \partial_\mu U^n \tau_3), \\ &= -\frac{i}{2n} \text{Tr} \left[\begin{pmatrix} A_1^* & A_2^* \\ -A_2 & A_1 \end{pmatrix} \partial_\mu \begin{pmatrix} A_1 & -A_2^* \\ A_2 & A_1^* \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \right], \\ &= -\frac{i}{2n} \text{Tr} \left[\begin{pmatrix} A_1^* & A_2^* \\ -A_2 & A_1 \end{pmatrix} \begin{pmatrix} \partial_\mu A_1 & \partial_\mu A_2^* \\ \partial_\mu A_2 & -\partial_\mu A_1^* \end{pmatrix} \right], \\ &= -\frac{i}{2n} \text{Tr} \begin{bmatrix} A_1^* \partial_\mu A_1 + A_2^* \partial_\mu A_2 & \\ & -A_2 \partial_\mu A_2^* - A_1 \partial_\mu A_1^* \end{bmatrix}, \\ &= -\frac{i}{2n} [\bar{A}^i \partial_\mu A_i - (\partial_\mu \bar{A}^i) A_i], \end{aligned}$$

$$\Rightarrow \mathbf{V}_\mu(\mathbf{r}) = -\frac{\mathbf{i}}{2\mathbf{n}} \bar{\mathbf{A}}^{\mathbf{i}} \vec{\partial} \mathbf{A}_{\mathbf{i}}. \quad (4.16)$$

With matrix $U^n(r)$ is defined in (4.11) and the gauge field is defined in (4.16), the supersymmetric Lagrangian density (4.3) is then

$$\mathcal{L}_n = -\frac{f_\pi^2}{8n^2} \bar{D}_\mu \bar{A} D^\mu A - \frac{1}{16e^2 n^2} F_{\mu\nu}^2(V), \quad (4.17)$$

with

$$F_{\mu\nu}(V) = \partial_\mu V_\nu - \partial_\nu V_\mu. \quad (4.18)$$

Really, we have

+ **The first term**

$$\begin{aligned} \mathcal{L}_n^1 &= -\frac{1}{16n^2} f_\pi^2 Tr \left[D^\mu \begin{pmatrix} A_1^* & A_2^* \\ -A_2 & A_1 \end{pmatrix} D_\mu \begin{pmatrix} A_1 & -A_2^* \\ A_2 & A_1^* \end{pmatrix} \right], \\ &= -\frac{1}{16n^2} f_\pi^2 Tr \left(\begin{array}{cc} D^\mu A_1^* D_\mu A_1 + D^\mu A_2^* D_\mu A_2 & \\ & D^\mu A_2 D_\mu A_2^* + D^\mu A_1 D_\mu A_1^* \end{array} \right), \\ &\Rightarrow \mathcal{L}_{\mathbf{n}}^1 = -\frac{1}{8\mathbf{n}^2} \mathbf{f}_\pi^2 \mathbf{D}^\mu \bar{\mathbf{A}}^{\mathbf{i}} \mathbf{D}_\mu \mathbf{A}_{\mathbf{i}}. \end{aligned} \quad (4.19)$$

+ **The second term**

$$\begin{aligned} &U^{-n} D_\mu U^n U^{-n} D_\nu U^n - U^{-n} D_\nu U^n U^{-n} D_\mu U^n, \\ &= U^{-n} [(\partial_\mu - iV_\mu \tau_3) U^n] U^{-n} [(\partial_\nu - iV_\nu \tau_3) U^n] - \\ &\quad - U^{-n} [(\partial_\nu - iV_\nu \tau_3) U^n] U^{-n} [(\partial_\mu - iV_\mu \tau_3) U^n], \\ &= (U^{-n} \partial_\mu U^n - iU^{-n} V_\mu \tau_3 U^n) (U^{-n} \partial_\nu U^n - iU^{-n} V_\nu \tau_3 U^n) - \\ &\quad - (U^{-n} \partial_\nu U^n - iU^{-n} V_\nu \tau_3 U^n) (U^{-n} \partial_\mu U^n - iU^{-n} V_\mu \tau_3 U^n), \\ &= U^{-n} \partial_\mu U^n U^{-n} \partial_\nu U^n - U^{-n} \partial_\nu U^n U^{-n} \partial_\mu U^n. \end{aligned} \quad (4.20)$$

Using

$$U^{-n} U^n = 1 \Rightarrow \partial_\mu U^{-n} U^n + U^{-n} \partial_\mu U^n = 0, \quad (4.21)$$

(4.20) is now

$$(4.20) = -\partial_\mu U^{-n} \partial_\nu U^n + \partial_\nu U^{-n} \partial_\mu U^n. \quad (4.22)$$

The second term is now

$$\Rightarrow \mathcal{L}_n^2 = \frac{1}{32e^2 n^4} \text{Tr} \left[(\partial_\mu U^{-n} \partial_\nu U^n - \partial_\nu U^{-n} \partial_\mu U^n)^2 \right]. \quad (4.23)$$

We define

$$F_{\mu\nu}(V) \equiv \partial_\mu V_\nu - \partial_\nu V_\mu, \quad (4.24)$$

$$= -\frac{i}{2n} \text{Tr} \left[\partial_\mu (U^{-n} \partial_\nu U^n \tau_3) - \partial_\nu (U^{-n} \partial_\mu U^n \tau_3) \right], \quad (4.25)$$

$$= -\frac{i}{2n} \text{Tr} \left[\partial_\mu U^{-n} \partial_\nu U^n \tau_3 - \partial_\nu U^{-n} \partial_\mu U^n \tau_3 \right]. \quad (4.26)$$

$$\Rightarrow F_{\mu\nu}^2 = \frac{1}{2n^2} \text{Tr} \left[(\partial_\mu U^{-n} \partial_\nu U^n - \partial_\nu U^{-n} \partial_\mu U^n)^2 \right]. \quad (4.27)$$

From (4.23) and (4.27), we have

$$\mathcal{L}_n^2 = \frac{1}{16e^2 n^2} \mathbf{F}_{\mu\nu}^2. \quad (4.28)$$

To supersymmetrise this model, we extend A_i to chiral scalar multiples $(A_i, \psi_{\alpha i}, F_i)$ ($i, \alpha = 1, 2$) and the vector $V_\mu(r)$ to real vector multiples $(V_\mu, \lambda_\alpha, D)$. Here, the fields F_i are complex scalars, D is real scalar, $\psi_{\alpha i}$, λ_α are Majorana two-component spinors. $\psi_{\alpha i}$ corresponds to a left-handed chiral spinor, $\bar{\psi}^{\alpha i} = (\psi_i^\alpha)^*$ corresponds to a right-handed one. The supersymmetric Lagrangian density is given by

$$\mathcal{L}_{susy} = \frac{f_\pi^2}{8n^2} \left[-D^\mu \bar{A}^i D_\mu A_i - \frac{1}{2} i \bar{\psi}^{\dot{\alpha} i} (\sigma_\mu)_{\alpha \dot{\alpha}} \vec{D}^\mu \psi_i^\alpha + \bar{F}^i F_i - \right. \quad (4.29)$$

$$\left. -i \bar{A}^i \lambda^\alpha \psi_{\alpha i} + i A_i \bar{\lambda}^{\dot{\alpha}} \bar{\psi}_{\dot{\alpha}}^i + D (\bar{A}^i A_i - 1) \right] + \quad (4.30)$$

$$+ \frac{1}{8e^2 n^2} \left[-\frac{1}{2} F_{\mu\nu}^2 - i \bar{\lambda}^{\dot{\alpha}} (\sigma^\mu)_{\dot{\alpha}}^\alpha \partial_\mu \lambda_\alpha + D^2 \right], \quad (4.31)$$

is invariant under supersymmetric transformations

$$\delta A_i = -\varepsilon^\alpha \psi_{\alpha i}, \quad (4.32)$$

$$\delta \psi_{\alpha i} = -i \bar{\varepsilon}^{\dot{\alpha}} (\sigma^\mu)_{\alpha \dot{\alpha}} D_\mu A_i + \varepsilon_\alpha F_i, \quad (4.33)$$

$$\delta F_i = -i \bar{\varepsilon}^{\dot{\alpha}} (\sigma^\mu)_{\dot{\alpha}}^\alpha D_\mu \psi_{\alpha i} - i \bar{\varepsilon}^{\dot{\alpha}} A_i \bar{\lambda}_{\dot{\alpha}}, \quad (4.34)$$

$$\delta V_\mu = -\frac{1}{2} i (\sigma_\mu)^{\alpha \dot{\alpha}} (\bar{\varepsilon}_{\dot{\alpha}} \lambda_\alpha + \varepsilon_\alpha \bar{\lambda}_{\dot{\alpha}}), \quad (4.35)$$

$$\delta \lambda_\alpha = \varepsilon^\beta (\sigma^{\mu\nu})_{\beta\alpha} F_{\mu\nu} + i \varepsilon_\alpha D, \quad (4.36)$$

$$\delta D = \frac{1}{2} (\sigma^\mu)_{\alpha \dot{\alpha}} \partial_\mu (\bar{\varepsilon}^{\dot{\alpha}} \lambda^\alpha - \varepsilon^\alpha \bar{\lambda}^{\dot{\alpha}}). \quad (4.37)$$

The field equation and their supersymmetric transformations lead to the following constraints with

$$\bar{A}^i A_i = 0, \quad (4.38)$$

$$\bar{A}^i \psi_{\alpha i} = 0, \quad (4.39)$$

$$\bar{A}^i F_i = 0 \quad (4.40)$$

and following algebraic expressions for

$$V_\mu = -\frac{1}{2} \left\{ i \bar{A}^i \vec{\partial}_\mu A_i + (\sigma_\mu)^{\alpha\dot{\alpha}} \bar{\psi}_{\dot{\alpha}}^i \psi_{\alpha i} \right\}, \quad (4.41)$$

$$\lambda_\alpha = -i \left\{ \bar{F}^i \psi_{\alpha i} + i (\sigma^\mu)_{\alpha\dot{\alpha}} (D_\mu A_i) \bar{\psi}^{\dot{\alpha} i} \right\}, \quad (4.42)$$

$$D = D^\mu \bar{A}^i D_\mu A_i + \frac{1}{2} i \bar{\psi}^{\dot{\alpha} i} (\sigma^\mu)_{\alpha\dot{\alpha}} \left(\vec{D}_\mu \psi_i^\alpha \right) - \bar{F}^i F_i. \quad (4.43)$$

Setting $\psi_\alpha = F_i = 0$, then

$$\mathcal{L}_{susy} = -\frac{f_\pi^2}{8n^2} \bar{D}^\mu \bar{A} D_\mu A + \frac{1}{8e^2 n^2} \left[-\frac{1}{2} F_{\mu\nu}^2 + (\bar{D}^\mu \bar{A} D_\mu A)^2 \right]. \quad (4.44)$$

The second term is fourth-order in derivatives. However, there is other possible fourth-order term

$$\square \bar{A} \square A - (\bar{D}^\mu \bar{A} D_\mu A)^2, \quad (4.45)$$

where $\square \equiv D^\mu D_\mu$ is the gauge covariant d'Alembertian.

Thus, we may let it to the Lagrangian density

$$\mathcal{L}_{susy} = -\frac{f_\pi^2}{8n^2} \bar{D}^\mu \bar{A} D_\mu A + \frac{1}{8e^2 n^2} \left[\alpha \left\{ -\frac{1}{2} F_{\mu\nu}^2 + (\bar{D}^\mu \bar{A} D_\mu A)^2 \right\} + \right. \quad (4.46)$$

$$\left. + \beta \left\{ \square \bar{A} \square A - (\bar{D}^\mu \bar{A} D_\mu A)^2 \right\} \right]. \quad (4.47)$$

Now, we recall the hedgehog ansatz

$$U^n(r) = \cos nf(r) + i \vec{\tau} \frac{\vec{r}}{r} \sin nf(r). \quad (4.48)$$

From (4.11), we have

$$A_1 = \cos nf(r) + i \cos \theta \sin nf(r), \quad (4.49)$$

$$A_2 = i e^{i\varphi} \sin \theta \sin nf(r). \quad (4.50)$$

Inserting the hedgehog ansatz into the supersymmetric Lagrangian density, we obtain the supersymmetric static energy

$$E = 4\pi \frac{f_\pi}{e} \int dx x^2 \left\{ \frac{1}{12} \left(f'^2 + \frac{2 \sin^2 nf}{n^2 x^2} \right) + \left(\frac{\alpha + \beta}{15} \right) \left(f'^2 - \frac{\sin^2 nf}{n^2 x^2} \right) + \right. \quad (4.51)$$

$$\left. + \frac{\beta}{12} \left(f'' + \frac{2f'}{nx} - \frac{\sin 2nf}{n^2 x^2} \right) \right\}. \quad (4.52)$$

The Euler-Lagrange equation takes the field equation

$$-x^2 f'' - 2xf' + \frac{\sin 2nf}{n} + \frac{4(\alpha + \beta)}{5} \left[\frac{2f'' \sin^2 nf}{n^2} - 6x^2 f'^2 f'' - 4xf'^3 + \right. \quad (4.53)$$

$$\left. + \frac{f'^2 \sin 2nf}{n} + \frac{\sin^2 nf \sin 2nf}{n^3 x^2} \right] + \beta \left[x^2 f^{(4)} + \frac{4xf^{(3)}}{n} - \frac{4f'' \cos 2nf}{n} + \right. \quad (4.54)$$

$$\left. + \frac{4f'^2 \sin 2nf}{n^2} - \frac{4 \sin^2 nf \sin 2nf}{n^3 x^2} \right] = 0, \quad (4.55)$$

with $x = ef_\pi r$ is the dimensionless variable and boundary conditions

$$f(0) = \pi, \quad (4.56)$$

$$f(\infty) = 0. \quad (4.57)$$




Figure 4.1: Plots of supersymmetric numerical solution of $f(x)$ for a case of $\alpha = 1, \beta = 0, n = 2, 3$ [26].

Chapter 5

Appendix

5.1 Some notations for Chapter 2 and 3

There are some notations to calculations in chapter 2 and 3

$$\frac{\partial}{\partial x^\mu} \equiv \partial_\mu = (\partial_0, \partial_i) = \left(\frac{\partial}{\partial t}, \vec{\nabla} \right); \quad (5.1)$$

$$\frac{\partial}{\partial x_\mu} \equiv \partial^\mu = (\partial^0, \partial^i) = g^{\mu\nu} \partial_\nu = \text{diag}(1, -1, -1, -1) \partial_\nu = \left(\frac{\partial}{\partial t}, -\vec{\nabla} \right); \quad (5.2)$$

$$\partial_i F(r) = \frac{\partial F(r)}{\partial x^i} = \frac{\partial F}{\partial r} \frac{\partial r}{\partial x^i} = F' \frac{x^i}{r}. \quad (5.3)$$

$$U_o^n(r) = \exp[i\vec{\tau} \cdot \hat{r} \{nF(r)\}] = \cos\{nF(r)\} + i\vec{\tau} \cdot \hat{r} \sin\{nF(r)\}; \quad (5.4)$$

$$U_o^{-n}(r) = \exp[-i\vec{\tau} \cdot \hat{r} \{nF(r)\}] = \cos\{nF(r)\} - i\vec{\tau} \cdot \hat{r} \sin\{nF(r)\}; \quad (5.5)$$

Some identities of Pauli matrices are

$$\tau_i \tau_j = i\varepsilon_{ijk} \tau_k; \quad (5.6)$$

$$(\tau_i)^2 = 1; \quad (5.7)$$

$$\tau_i \tau_j = -\tau_j \tau_i; \quad (5.8)$$

$$\text{Tr}(\tau_i^2) = 2; \quad (5.9)$$

$$\text{Tr}(\tau_i) = 0; \quad (5.10)$$

We will define

$$\begin{aligned} \partial_\mu U_o^n(r) &= \partial_\mu (\cos nF) + i\vec{\tau} \partial_\mu (\hat{r} \sin nF), \\ &= -n\partial_i F \sin nF + \frac{i\vec{\tau}}{r} \partial_i \vec{r} \sin nF + i\vec{\tau} \vec{r} \partial_i \left(\frac{1}{r} \right) \sin nF + in\vec{\tau} \frac{\vec{r}}{r} \partial_i F \cos nF, \end{aligned}$$

$$= -n \frac{F' x^i}{r} \sin nF + \frac{i\tau_i \vec{e}^i}{r} \sin nF - \frac{i\vec{\tau}\vec{r}x^i}{r^3} \sin nF + in \frac{\vec{\tau}\vec{r}F' x^i}{r^2} \cos nF; \quad (5.11)$$

$$\begin{aligned} \partial_\mu U_o^{-n}(r) &= \partial_\mu (\cos nF) - i\vec{\tau}\partial_\mu (\hat{r} \sin nF), \\ &= -n \frac{F' x^i}{r} \sin nF - \frac{i\tau_i \vec{e}^i}{r} \sin nF + \frac{i\vec{\tau}\vec{r}x^i}{r^3} \sin nF - in \frac{\vec{\tau}\vec{r}F' x^i}{r^2} \cos nF; \end{aligned} \quad (5.12)$$

$$\begin{aligned} \partial^\mu U_o^n(r) &= \partial^\mu (\cos nF) + i\vec{\tau}\partial^\mu (\hat{r} \sin nF), \\ &= n \frac{F' x_i}{r} \sin nF - \frac{i\tau_i \vec{e}_i}{r} \sin nF + \frac{i\vec{\tau}\vec{r}x_i}{r^3} \sin nF - in \frac{\vec{\tau}\vec{r}F' x_i}{r^2} \cos nF; \end{aligned} \quad (5.13)$$

The first term of the Lagrangian density (2.2) is defined

$$\begin{aligned} &\Rightarrow \mathcal{L}_n^1 = Tr (\partial_\mu U_o^{-n} \partial^\mu U_o^n) \\ &= Tr \left\{ \left(-n \frac{F' x^i}{r} \sin nF - \frac{i\tau_i \vec{e}^i}{r} \sin nF + \frac{i\vec{\tau}\vec{r}x^i}{r^3} \sin nF - in \frac{\vec{\tau}\vec{r}F' x^i}{r^2} \cos nF \right) \times \right. \\ &\quad \times \left. \left(n \frac{F' x_i}{r} \sin nF - \frac{i\tau_i \vec{e}_i}{r} \sin nF + \frac{i\vec{\tau}\vec{r}x_i}{r^3} \sin nF - in \frac{\vec{\tau}\vec{r}F' x_i}{r^2} \cos nF \right) \right\} \\ &= Tr \left\{ -n^2 F'^2 \sin^2 nF + in F' \frac{\vec{\tau}\vec{r}}{r^2} \sin^2 nF - in F' \frac{\vec{\tau}\vec{r}}{r^2} \sin^2 nF + \right. \\ &\quad + in^2 F'^2 \frac{\vec{\tau}\vec{r}}{2r} \sin 2nF - in F' \frac{\vec{\tau}\vec{r}}{r^2} \sin^2 nF - 3 \frac{\sin^2 nF}{r^2} + \frac{\sin^2 nF}{r^2} - \\ &\quad - n \frac{F'}{2r} \sin 2nF + in F' \frac{\vec{\tau}\vec{r}}{r^2} \sin^2 nF + \frac{\sin^2 nF}{r^2} - \frac{\sin^2 nF}{r^2} + n \frac{F'}{2r} \sin 2nF - \\ &\quad \left. - in^2 F'^2 \frac{\vec{\tau}\vec{r}}{2r} \sin 2nF - n \frac{F'}{2r} \sin 2nF + n \frac{F'}{2r} \sin 2nF - n^2 F'^2 \cos^2 nF \right\} \\ &= Tr \left(-n^2 F'^2 - 2 \frac{\sin^2 nF}{r^2} \right) \\ &\Rightarrow \mathcal{L}_n^1 = -2n^2 \left(\frac{dF}{dr} \right)^2 - 4 \frac{\sin^2 nF}{r^2}. \end{aligned} \quad (5.14)$$

To defined the second term of the Lagrangian density (2.2), we will go to the intermediary step using the unitary condition of U_o^n

$$U_o^{-n} U_o^n = 1 \Rightarrow \partial_\mu U_o^{-n} U_o^n + U_o^{-n} \partial_\mu U_o^n = 0. \quad (5.15)$$

The second term of Lagrangian density (2.2) is now

$$\Rightarrow \mathcal{L}_n^2 = Tr \left[(\partial_\mu U_o^{-n} \partial_\nu U_o^n - \partial_\nu U_o^{-n} \partial_\mu U_o^n)^2 \right]. \quad (5.16)$$

Note (5.11), (5.12) we can define

$$\mathcal{L}_n^2 = 32n^2 \frac{\sin^2 nF}{r^2} \left(\frac{dF}{dr} \right)^2 + 16 \frac{\sin^4 nF}{r^4}. \quad (5.17)$$

5.2 Some notations for Chapter 4

Dotted and undotted spinor indices run from 1 to 2 and are denoted by the early letters of the English alphabet. Spinor indices are raised and lowered by the ε tensors

$$\varepsilon^{ab} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad (5.18)$$

$$\varepsilon_{ab} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}. \quad (5.19)$$

For example

$$\psi^a = \varepsilon^{ab} \psi_b, \quad (5.20)$$

$$\chi^a = \varepsilon_{ab} \chi^b. \quad (5.21)$$

The product $\psi\chi$ is

$$\psi\chi = \psi^a \chi_a = -\psi_a \chi^a = \chi^a \psi_a = \chi\psi. \quad (5.22)$$

In the special case

$$\psi^a \psi^b = -\frac{1}{2} \psi\psi \varepsilon^{ab}. \quad (5.23)$$

The hermitian conjugates of the spinors ψ and χ are $\bar{\psi}$ and $\bar{\chi}$

$$\bar{\psi}^{\dot{a}} = (\psi^a)^\dagger, \quad (5.24)$$

$$\bar{\chi} = (\chi_a)^\dagger. \quad (5.25)$$

Their dotted indices are raised and lowered by the tensors $\varepsilon^{\dot{a}\dot{b}}$ and $\varepsilon_{\dot{a}\dot{b}}$ which are equal to their undotted counterparts ε^{ab} and ε_{ab} . The product of them is

$$\bar{\psi}\bar{\chi} = \bar{\psi}_{\dot{a}} \bar{\chi}^{\dot{a}} = -\bar{\psi}^{\dot{a}} \bar{\chi}_{\dot{a}} = \bar{\chi}_{\dot{a}} \bar{\psi}^{\dot{a}} = \bar{\chi}\bar{\psi}. \quad (5.26)$$

In the special case

$$\bar{\psi}^{\dot{a}} \bar{\psi}^{\dot{b}} = \frac{1}{2} \bar{\psi}\bar{\psi} \varepsilon^{\dot{a}\dot{b}}. \quad (5.27)$$

The hermitian conjugates of the product $\chi\psi$ is

$$(\chi\psi)^\dagger = (\chi^a \psi_a)^\dagger = \bar{\psi}_{\dot{a}} \bar{\chi}^{\dot{a}} = \bar{\psi}\bar{\chi} = \bar{\chi}\bar{\psi}. \quad (5.28)$$

The Pauli matrices σ_{ab}^m are

$$\begin{aligned}\sigma^0 &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \sigma^1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \\ \sigma^2 &= \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \sigma^3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.\end{aligned}\quad (5.29)$$

Other useful identities are

$$\chi \sigma^n \bar{\psi} = -\bar{\psi} \bar{\sigma}^n \chi, \quad (5.30)$$

$$\chi \sigma^m \bar{\sigma}^n \psi = \psi \sigma^n \bar{\sigma}^m \chi, \quad (5.31)$$

$$(\chi \sigma^m \bar{\psi})^\dagger = \psi \sigma^m \bar{\chi}, \quad (5.32)$$

$$(\chi \sigma^m \bar{\sigma}^n \psi)^\dagger = \bar{\psi} \bar{\sigma}^n \sigma^m \bar{\chi}. \quad (5.33)$$

Using these above notations, we can define (from...)

$$\delta \bar{A}_i = -\bar{\psi}_{\dot{\alpha}i} \bar{\varepsilon}^{\dot{\alpha}}, \quad (5.34)$$

$$\delta \bar{\psi}_{\dot{\alpha}i} = i (\sigma^\mu)_{\alpha\dot{\alpha}} \varepsilon^\alpha D_\mu \bar{A}_i + \bar{\varepsilon}_{\dot{\alpha}} \bar{F}_i, \quad (5.35)$$

$$\delta \bar{F}_i = i D_\mu \bar{\psi}_{\dot{\alpha}i} (\sigma^\mu)_{\dot{\alpha}}^\alpha \varepsilon^\alpha + i \bar{A}_i \lambda_\alpha \varepsilon^\alpha, \quad (5.36)$$

$$\delta \bar{V}_\mu = \frac{1}{2} i (\bar{\lambda}_{\dot{\alpha}} \varepsilon_\alpha + \lambda_\alpha \bar{\varepsilon}_{\dot{\alpha}}) (\sigma_\mu)^{\alpha\dot{\alpha}}, \quad (5.37)$$

$$\delta \bar{\lambda}_{\dot{\alpha}} = (\sigma^{\mu\nu})_{\beta\alpha} \bar{\varepsilon}^{\dot{\beta}} F_{\mu\nu} - i \bar{\varepsilon}_{\dot{\alpha}} \bar{D}, \quad (5.38)$$

$$\delta \bar{D} = \frac{1}{2} \partial_\mu (\bar{\lambda}^{\dot{\alpha}} \varepsilon^\alpha - \lambda^\alpha \bar{\varepsilon}^{\dot{\alpha}}) (\sigma^\mu)_{\alpha\dot{\alpha}}. \quad (5.39)$$

Since δ and D_μ can permute together, $\delta \mathcal{L}_{susy}$ is defined

$$\begin{aligned}\delta \mathcal{L}_{susy} &= \frac{f_\pi^2}{8n^2} \left[-D^\mu (\delta \bar{A}^i) D_\mu A_i - D^\mu \bar{A}^i D_\mu (\delta A_i) - \frac{1}{2} i (\delta \bar{\psi}^{\dot{\alpha}i}) (\sigma_\mu)_{\alpha\dot{\alpha}} \bar{D}^{\mu} \psi_i^\alpha - \right. \\ &\quad \left. - \frac{1}{2} i \bar{\psi}^{\dot{\alpha}i} (\sigma_\mu)_{\alpha\dot{\alpha}} \bar{D}^{\mu} (\delta \psi_i^\alpha) + \delta \bar{F}^i F_i + \bar{F}^i \delta F_i - i (\delta \bar{A}^i) \lambda^\alpha \psi_{\alpha i} - i \bar{A}^i (\delta \lambda^\alpha) \psi_{\alpha i} - i \bar{A}^i \lambda^\alpha \delta \psi_{\alpha i} + \right. \\ &\quad \left. + i (\delta A_i) \bar{\lambda}^{\dot{\alpha}} \bar{\psi}_{\dot{\alpha}}^i + i A_i (\delta \bar{\lambda}^{\dot{\alpha}}) \bar{\psi}_{\dot{\alpha}}^i + i A_i \bar{\lambda}^{\dot{\alpha}} (\delta \bar{\psi}_{\dot{\alpha}}^i) + (\delta D) (\bar{A}^i A_i - 1) + D (\delta \bar{A}^i) A_i + D \bar{A}^i (\delta A_i) \right] + \\ &\quad + \frac{1}{8e^2 n^4} \left[-F_{\mu\nu} \delta F_{\mu\nu} - i (\delta \bar{\lambda}^{\dot{\alpha}}) (\sigma^\mu)_{\dot{\alpha}}^\alpha \partial_\mu \lambda_\alpha - i \bar{\lambda}^{\dot{\alpha}} (\sigma^\mu)_{\dot{\alpha}}^\alpha \partial_\mu (\delta \lambda_\alpha) + 2D \delta D \right]. \quad (5.40)\end{aligned}$$

Bibliography

- [1] T. H. R. Skyrme, *Nucl. Phys.*, **31** (1962) 566.
- [2] G. S. Adkins, C. R. Nappi and E. Witten, *Nucl. Phys.*, **B228** (1983) 552. G. S. Adkins, *Nucl. Phys.*, **B249** (1985) 507. G.S. Adkins, C. R. Nappi, *Nucl. Phys.*, **B233** (1984) 109. E. Witten, *Nucl. Phys.*, **B223** (1983) 422. E. Witten, *Nucl. Phys.*, **B223** (1983) 433.
- [3] N. A. Viet, P. T. Tuyen, *J. Phys.*, **G15** (July 1989) 937.
- [4] H. Y. Cheung, Feza Gursey, *Modern Physics Letters A*, **Vol.5 No.21** (1990) 1685
- [5] E. A. Bergshoeff, R. I. Nepomachie, H. J. Schnitzer, *Nucl. Phys.*, **B249** (1985) 93
- [6] N. Shiiki, N. Sawado, S.Oryu, hep-th/0603069 (2006)
- [7] A. Acus, hep-ph/9901240 (1999)
- [8] C. Houghton, S. Magee, hep-th/0602227 (2006)
- [9] J. Wess, B. Zumino, *Nucl. Phys.*, **B70** (1974) 39. J. Wess, B. Zumino, *Nucl. Phys.*, **B78** (1974) 1.
- [10] A. Salam, J. Strathdee, *Nucl. Phys.*, **B76** (1974) 477. A. Salam, J. Strathdee, *Nucl. Phys.*, **B80** (1974) 499.
- [11] M. F. Sohnius, *Physics Report*, **128, Nos 2 and 3**, (1985), 39 - 204
- [12] P. T. Tuyen, *Quantum Mechanics, in Vietnamese*, College of Science, VNU, (2004)
- [13] P. T. Tuyen, *Theory of Fundamental Particles, in Vietnamese*, College of Science, VNU, (2006)
- [14] P. T. Tuyen, *Theory of Group, in Vietnamese*, College of Science, VNU, (2006)
- [15] L. Ryder, *Quantum Field Theory, second edition*, Cambridge University Press, (1996)
- [16] S. Weinberg, *The Quantum Theory of Fields*, **vol I, II, III**, Cambridge University Press, (1995)
- [17] M. E. Peskin, D. V. Shroeder, *An Introduction to Quantum Field Theory*, Westview Press, (1995)
- [18] H. Georgi, *Lie Algebras in Particle Physics, second edition*, Perseus Books, (1999)
- [19] W. Greiner, J. Reinhardt, *Field Quantization*, Springer, (1996)
- [20] A. Bilal, hep-th/0101055 (2001)
- [21] K. Cahill, hep-ph/9907295 (1999)
- [22] D. Q. Tuan, P. T. Tuyen, *J. Science. (Math - Phys)*, VNU, **T.XXII, No 2AP** (2006) 194

- [23] D. Q. Tuan, P. T. Tuyen, **Phenomenal models of Baryon**, *Talk in National Conference on Theoretical Physics 31*, Cua Lo, Nghe An, 2006 (no-published)
- [24] D. Q. Tuan, P. T. Tuyen, **The composite Skyrme model with the mass of pion**, (no-published) 2006
- [25] D. Q. Tuan, P. T. Tuyen, **The composite Skyrme model with N=4**, (no-published) 2006
- [26] D. Q. Tuan, P. T. Tuyen, **The supersymmetric composite skyrmion**, (no-published) 2007
- [27] Nikolaev V.A. , Tkachev O.G. , hep-ph/0109192 (2001)
- [28] O.V.Pavlovskii, hep-th/0312343 (2003)
- [29] R. A. Battye, P. M. Sutcliffe, hep-th/0602220 (2006)
- [30] D. Q. Tuan, P. T. Tuyen, **The formalism of gravitational composite Skyrme model**, *Talk in National Conference on Theoretical Physics 32*, Nha Trang, Khanh Hoa, 2007 (no-published)